

Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname

\_\_\_\_\_

Forename(s)

\_\_\_\_\_

Candidate signature

\_\_\_\_\_

I declare this is my own work.

# GCSE CHEMISTRY

# H

Higher Tier Paper 1

Thursday 14 May 2020

Morning

Time allowed: 1 hour 45 minutes

## Materials

For this paper you must have:

- a ruler
- a calculator
- the periodic table (enclosed).

## Instructions

- Use black ink or black ball-point pen.
- Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do all rough work in this book. Cross through any work you do not want to be marked.
- In all calculations, show clearly how you work out your answer.

## Information

- The maximum mark for this paper is 100.
- The marks for questions are shown in brackets.
- You are expected to use a calculator where appropriate.
- You are reminded of the need for good English and clear presentation in your answers.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
<b>TOTAL</b>	



0 1

This question is about structure and bonding.

0 1 . 1

Which **two** substances have intermolecular forces between particles?**[2 marks]**Tick (✓) **two** boxes.

Diamond

Magnesium

Poly(ethene)

Sodium chloride




Water

0 1 . 2

Table 1 shows the structures of three compounds.

Table 1

Diagrams not to scale

Compound	Structure
Carbon dioxide	 <p>Key</p> <ul style="list-style-type: none"> <li><span style="display: inline-block; width: 10px; height: 10px; background-color: black; border-radius: 50%;"></span> O</li> <li><span style="display: inline-block; width: 10px; height: 10px; background-color: grey; border-radius: 50%;"></span> C</li> </ul>
Magnesium oxide	 <p>Key</p> <ul style="list-style-type: none"> <li><span style="display: inline-block; width: 10px; height: 10px; background-color: black; border-radius: 50%;"></span> O<sup>2-</sup></li> <li><span style="display: inline-block; width: 10px; height: 10px; background-color: grey; border-radius: 50%;"></span> Mg<sup>2+</sup></li> </ul>
Silicon dioxide	 <p>Key</p> <ul style="list-style-type: none"> <li><span style="display: inline-block; width: 10px; height: 10px; background-color: black; border-radius: 50%;"></span> O</li> <li><span style="display: inline-block; width: 10px; height: 10px; background-color: grey; border-radius: 50%;"></span> Si</li> </ul>





**0 2**

This question is about metals and the reactivity series.

**0 2 . 1**Which **two** statements are properties of most transition metals?**[2 marks]**Tick (✓) **two** boxes.

They are soft metals.

They form colourless compounds.

They form ions with different charges.

They have high melting points.

They have low densities.

**0 2 . 2**

A student added copper metal to colourless silver nitrate solution.

The student observed:

- pale grey crystals forming
- the solution turning blue.

Explain how these observations show that silver is less reactive than copper.

**[3 marks]**

---

---

---

---

---

---

---





**0 2 . 4** Metal **M** has two isotopes.

**Table 2** shows the mass numbers and percentage abundances of the isotopes.

**Table 2**

Mass number	Percentage abundance (%)
203	30
205	70

Calculate the relative atomic mass ( $A_r$ ) of metal **M**.

Give your answer to 1 decimal place.

**[2 marks]**

---

---

---

---

Relative atomic mass (1 decimal place) = \_\_\_\_\_

**11**



**Turn over for the next question**

*Do not write  
outside the  
box*

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**

**Turn over ►**




**0 3**

This question is about silver iodide.

Silver iodide is produced in the reaction between silver nitrate solution and sodium iodide solution.

The equation for the reaction is:

**0 3 . 1**

A student investigated the law of conservation of mass.

This is the method used.

1. Pour silver nitrate solution into a beaker labelled **A**.
2. Pour sodium iodide solution into a beaker labelled **B**.
3. Measure the masses of both beakers and their contents.
4. Pour the solution from beaker **B** into beaker **A**.
5. Measure the masses of both beakers and their contents again.

**Table 3** shows the student's results.

**Table 3**

	Mass before mixing in g	Mass after mixing in g
Beaker <b>A</b> and contents	78.26	108.22
Beaker <b>B</b> and contents	78.50	48.54

Explain how the results demonstrate the law of conservation of mass.

You should use data from **Table 3** in your answer.

**[2 marks]**

---

---

---

---





0 3 . 2

Suggest how the student could separate the insoluble silver iodide from the mixture at the end of the reaction.

[1 mark]

---

---

The student purified the separated silver iodide.

This is the method used.

1. Rinse the silver iodide with distilled water.
2. Warm the silver iodide.

0 3 . 3

Suggest **one** impurity that was removed by rinsing with water.

[1 mark]

---

---

0 3 . 4

Suggest why the student warmed the silver iodide.

[1 mark]

---

---

Question 3 continues on the next page


Turn over ►



03.5

Calculate the percentage atom economy for the production of silver iodide in this reaction.

The equation for the reaction is:



Give your answer to 3 significant figures.

Relative formula masses ( $M_r$ ):  $\text{AgNO}_3 = 170$   $\text{NaI} = 150$   $\text{AgI} = 235$   $\text{NaNO}_3 = 85$

[4 marks]

---

---

---

---

---

---

---

---

---

---

Percentage atom economy (3 significant figures) = \_\_\_\_\_ %

03.6

Give **one** reason why reactions with a high atom economy are used in industry.

[1 mark]

---

---

10



**Turn over for the next question**

*Do not write  
outside the  
box*

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**

**Turn over ►**



0 4

This question is about electrolysis.

A student investigated the electrolysis of copper chromate solution.


Copper chromate solution is green.

Copper chromate contains:

- blue coloured  $\text{Cu}^{2+}$  ions
- yellow coloured  $\text{CrO}_4^{2-}$  ions.

**Figure 1** shows the apparatus used.

**Figure 1**



The student switched the power supply on.

The student observed the changes at each electrode.

**Table 4** shows the student's observations.

**Table 4**

Changes at positive electrode	Changes at negative electrode
Solution turned yellow	Solution turned blue
Bubbles formed at the electrode	Solid formed on the electrode



0 4 . 1

Explain why the colour changed at the positive electrode.

**[2 marks]**

---

---

---

---

0 4 . 2

The gas produced at the positive electrode was oxygen.

The oxygen was produced from hydroxide ions.

Name the substance in the solution that provides the hydroxide ions.

**[1 mark]**

---

0 4 . 3

Describe how the solid forms at the negative electrode.

**[3 marks]**

---

---

---

---

---

---

---

0 4 . 4

The student repeated the investigation using potassium iodide solution instead of copper chromate solution.

Name the product at each electrode when potassium iodide solution is electrolysed.

**[2 marks]**

Negative electrode \_\_\_\_\_

Positive electrode \_\_\_\_\_


8

**Turn over ►**

0 5

This question is about the development of scientific theories.

**Figure 2** shows a timeline of some important steps in the development of the model of the atom.



0 5 . 1

The plum pudding model did not have a nucleus.

Describe **three** other differences between the nuclear model of the atom and the plum pudding model.

[3 marks]

- 1 \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_
- 2 \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_
- 3 \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_



**0 5 . 2** Niels Bohr adapted the nuclear model.

Describe the change that Bohr made to the nuclear model.

**[2 marks]**

---

---

---

---

**0 5 . 3** Mendeleev published his periodic table in 1869.

Mendeleev arranged the elements in order of atomic weight.

Mendeleev then reversed the order of some pairs of elements.

A student suggested Mendeleev's reason for reversing the order was to arrange the elements in order of atomic number.

Explain why the student's suggestion **cannot** be correct.

Use **Figure 2**.

**[2 marks]**

---

---

---

---

**0 5 . 4** Give the correct reason why Mendeleev reversed the order of some pairs of elements.  
**[1 mark]**

---

---







Magnesium displaces zinc from zinc sulfate solution.

**0 6 . 3** Complete the ionic equation for the reaction.

You should include state symbols.

**[2 marks]**



**0 6 . 4** Explain why the reaction between magnesium atoms and zinc ions is both oxidation and reduction.

**[2 marks]**

---

---

---

---

**9**

**Turn over for the next question**

**Turn over ►**




0 7

The reaction between hydrogen and oxygen releases energy.

0 7 . 1

A student drew a reaction profile for the reaction between hydrogen and oxygen.

**Figure 3** shows the student's reaction profile.



The student made **two** errors when drawing the reaction profile.

Describe the **two** errors.

**[2 marks]**

1 \_\_\_\_\_

2 \_\_\_\_\_

\_\_\_\_\_



**0 7 . 2** The reaction between hydrogen and oxygen in a hydrogen fuel cell is used to produce electricity.

Hydrogen fuel cells and rechargeable cells are used to power some cars.

Give **two** advantages of using hydrogen fuel cells instead of using rechargeable cells to power cars.

**[2 marks]**

1 \_\_\_\_\_

2 \_\_\_\_\_

**0 7 . 3** Reactions occur at the positive electrode and at the negative electrode in a hydrogen fuel cell.

Write a half equation for **one** of these reactions.

**[1 mark]**

\_\_\_\_\_

**Question 7 continues on the next page**


**Turn over ►**



**0 7 . 4** The three states of matter can be represented by a simple particle model.

**Figure 4** shows a simple particle model for hydrogen gas.

**Figure 4**



Give **two** limitations of this simple particle model for hydrogen gas.

**[2 marks]**

1 \_\_\_\_\_

2 \_\_\_\_\_

**0 7 . 5** The hydrogen gas needed to power a car for 400 km would occupy a large volume.

Suggest **one** way that this volume can be reduced.

**[1 mark]**

\_\_\_\_\_

\_\_\_\_\_



0 7 . 6

The energy needed for a car powered by a hydrogen fuel cell to travel 100 km is 58 megajoules (MJ).

The energy released when 1 mole of hydrogen gas reacts with oxygen is 290 kJ

The volume of 1 mole of a gas at room temperature and pressure is 24 dm<sup>3</sup>

Calculate the volume of hydrogen gas at room temperature and pressure needed for the car to travel 100 km

**[4 marks]**

---

---

---

---

---

---

---

---

---

---

Volume of hydrogen gas = \_\_\_\_\_ dm<sup>3</sup>

**12****Turn over for the next question****Turn over ►**

0 8

This question is about the halogens.

**Table 5** shows the melting points and boiling points of some halogens.**Table 5**

Element	Melting point in °C	Boiling point in °C
Fluorine	-220	-188
Chlorine	-101	-35
Bromine	-7	59

0 8 . 1

What is the state of bromine at 0 °C **and** at 100 °C?**[1 mark]**Tick (✓) **one** box.**State at 0 °C****State at 100 °C**

Gas

Gas

Gas

Liquid

Liquid

Gas

Liquid

Liquid

Solid

Gas

Solid

Liquid



**0 8 . 2** Explain the trend in boiling points of the halogens shown in **Table 5**.

**[4 marks]**

---

---

---

---

---

---

---

---

---

---

**0 8 . 3** Why is it **not** correct to say that the boiling point of a single bromine molecule is 59 °C?

**[1 mark]**

---

---


**Question 8 continues on the next page**

**Turn over ►**



Iron reacts with each of the halogens in their gaseous form.

**Figure 5** shows the apparatus used.



**0 8 . 4** Give **one** reason why this experiment should be done in a fume cupboard.

**[1 mark]**

---

---

**0 8 . 5** Explain why the reactivity of the halogens decreases going down the group.

**[3 marks]**

---

---

---

---

---

---






0 8 . 6

A teacher investigated the reaction of iron with chlorine using the apparatus in **Figure 5**.

The word equation for the reaction is:



The teacher weighed:

- the glass tube
- the glass tube and iron before the reaction
- the glass tube and iron chloride after the reaction.

**Table 6** shows the teacher's results.

**Table 6**

	Mass in g
Glass tube	51.56
Glass tube and iron	56.04
Glass tube and iron chloride	64.56

Calculate the simplest whole number ratio of:

moles of iron atoms : moles of chlorine atoms

Determine the balanced equation for the reaction.

Relative atomic masses ( $A_r$ ): Cl = 35.5 Fe = 56

**[6 marks]**

---



---



---



---



---



---



---

Moles of iron atoms : moles of chlorine atoms = \_\_\_\_\_ : \_\_\_\_\_

Equation for the reaction \_\_\_\_\_

16

Turn over ►



0 9

This question is about citric acid ( $C_6H_8O_7$ ).

Citric acid is a solid.

A student investigated the temperature change during the reaction between citric acid and sodium hydrogencarbonate solution.


This is the method used.

1. Pour  $25\text{ cm}^3$  of sodium hydrogencarbonate solution into a polystyrene cup.
2. Measure the temperature of the sodium hydrogencarbonate solution.
3. Add  $0.20\text{ g}$  of citric acid to the polystyrene cup.
4. Stir the solution.
5. Measure the temperature of the solution.
6. Repeat steps 3 to 5 until a total of  $2.00\text{ g}$  of citric acid has been added.

The student plotted the results on a graph.

**Figure 6** shows the student's graph.

**Figure 6**



0 9 . 1

**Figure 6** shows an anomalous point when 0.60 g of citric acid was added. This was caused by the student making an error.

The student correctly:

- measured the mass of the citric acid
- read the thermometer
- plotted the point.

Suggest **one** reason for the anomalous point.

[1 mark]

---

---

0 9 . 2

Explain the shape of the graph in terms of the energy transfers taking place.

You should use data from **Figure 6** in your answer.

[3 marks]

---

---

---

---

---

---

0 9 . 3

A second student repeated the investigation using a metal container instead of the polystyrene cup. The container and the cup were the same size and shape.

Sketch a line on **Figure 6** to show the second student's results until 1.00 g of citric acid had been added. The starting temperature of the solution was the same.

Explain your answer.

[3 marks]

---

---

---

---

Turn over ►



The student used a solution of citric acid to determine the concentration of a solution of sodium hydroxide by titration.

0 9 . 4

The student made  $250 \text{ cm}^3$  of a solution of citric acid of concentration  $0.0500 \text{ mol/dm}^3$ .  
Calculate the mass of citric acid ( $\text{C}_6\text{H}_8\text{O}_7$ ) required.

Relative atomic masses ( $A_r$ ): H = 1 C = 12 O = 16

[3 marks]

---

---

---

---

---

Mass = \_\_\_\_\_ g

This is part of the method the student used for the titration.

1. Measure  $25.0 \text{ cm}^3$  of the sodium hydroxide solution into a conical flask using a pipette.
2. Add a few drops of indicator to the flask.
3. Fill a burette with citric acid solution.

0 9 . 5

Describe how the student would complete the titration.

[3 marks]

---

---

---

---

---

---



**0 9 . 6** Give **two** reasons why a burette is used for the citric acid solution.


**[2 marks]**

1 \_\_\_\_\_

2 \_\_\_\_\_

**0 9 . 7** 13.3 cm<sup>3</sup> of 0.0500 mol/dm<sup>3</sup> citric acid solution was needed to neutralise 25.0 cm<sup>3</sup> of sodium hydroxide solution.

The equation for the reaction is:



Calculate the concentration of the sodium hydroxide solution in mol/dm<sup>3</sup>

**[3 marks]**

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

Concentration = \_\_\_\_\_ mol/dm<sup>3</sup>

**18**

**END OF QUESTIONS**



**There are no questions printed on this page**

*Do not write  
outside the  
box*

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**





